



# Liver resection for synchronous metastasis from adenocarcinoma of the pâncreas

Orlando J M Torres
Pancreas study group

## PANCREAS STUDY GROUP


Fabrizio Di Benedetto (Modena University, Italy)
Orlando J. Torres (Maranhão Federal University, Brazil)



• 33 participants, 50 cases from 11 centers:




## Gender






## Pancreatic tumor location



Preoperatively pancreatic tumor size (Mean ± SD)	29.35 ± 17.65
Preoperatively CA 19-9 (Mean ± SD)	826.9 ± 1951.69




## Diagnosis methodology






## Staging at diagnosis








## **Vascular infiltration**






## Liver metastasis



Maximum Size of liver metastasis (Mean ±SD)	15.27 ± 9.57



## **Extrahepatic metastasis**




Number of positive nodes (Mean ± SD)	3.44 ± 4.39
Number of active metastasis (Mean ±SD)	1.16 ± 0.91

**Pancreas Study Group** 




## Neoadjuvant chemotherapy






## **Neoadjuvant chemotherapy**

#### Neoadjuvant chemotherapy %




## Liver segments involved








## Pancreatic procedure











## Liver segments resected








#### **Vascular resection**






## **Complications Grade A**







## R Status





## Adjuvant chemo and radiotherapy






Number of metastasis after adjuvant therapy (Mean ± SD) 0.72 ± 1.71
---

**Pancreas Study Group** 




#### **Overall survival**






## Overall survival



**Pancreas Study Group** 




#### Disease-free survival





## Disease-free survival







Thanks!